

Table 2 Fungicide seed treatments for wheat and barley and diseases controlled

	Common Bunt wheat and Covered Smut barley	Loose Smut barley	Dwarf Bunt wheat	Pythium Seed rot	Dry Seed/seedling decay	Fusarium crown/root rot- soil borne	Fusarium root rot-caused by scabby seed	Common root rot	Bare patch (Rhizoctonia)
Allegiance FL & LS Apron XL LS				E	E				
Charter Charter PB	E	E			E*	E	Not tested	E	E
Captan Maneb Mancozeb Thiram	E			V	E		E		
Dividend Extreme Dividend XL RTA	E		E	E	E	E	E	E	E
Maxim					E	E	E	E	
Raxil Thiram	E	E		V	E	E	E	E	E
Raxil XT and MD	E	E		E	E	E	E	E	E
Vitavax RTU Vitavax Extra	E	E		V	E	V**	E**	V**	E

(**E**)=Effective disease control ()= disease not controlled (**V**)=may give some control (*****)= Charter PB more effective than Charter alone (******) Vitavax Extra more effective than Vitavax RTU.

Source: Small Grain Seed Treatment Guide 2007. Alan Dyer, marry Burrows, Bob Johnston and Cecil Tharp. Montana State University Extension, MT 199608 AG 3/07