

Use Directions Update



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Intrepid[®] 2F

EPA Reg. No. 62719-442

Control of Lepidoptera Larvae Infesting Highbush Blueberries

(Refer to product label for General Use Precautions, Mixing and Application instructions.)

Pests, Application Rates, Application Timing and Restrictions:

Pests	Application Rate (fl oz/acre)	Application Timing	Restrictions
Fruitworm: cherry cranberry	12 - 16 (0.16 - 0.25 lb ai/acre)	Apply at initiation of egg laying (full bloom). Make subsequent applications as necessary depending on pest pressure or sustained moth flight.	<ul style="list-style-type: none"> Do not apply more than 64 fl oz of Intrepid 2F (1 lb ai) per acre per calendar year. Do not make more than 3 applications per acre per calendar year. Re-treatment Interval: 7 days Pre-harvest Interval: Do not within 7 days of harvest.
Leafroller: oblique-banded redbanded variegated	10 - 16 (0.19 - 0.25 lb ai/acre)	<p>Oblique-banded leafroller - overwintering generation: make one or two applications from pink bud through bloom and petal fall to small larvae when threshold levels occur.</p> <p>Other leafrollers and oblique-banded leafroller summer generation: apply at early egg hatch for each generation before webbing and sheltering begins. Make subsequent applications as necessary depending on pest pressure or sustained moth flight.</p>	
spanworm green fruitworm		Apply when larvae are first detected, feeding damage appears or when infestations reach threshold levels as defined by cooperative extension service or other qualified professional authorities. Make subsequent applications as necessary depending on pest pressure or sustained moth flight.	
armyworm cutworm	8 - 16 (0.12 - 0.25 lb ai/acre)	Apply when first signs of feeding damage appear or when infestations reach threshold levels as defined by cooperative extension service or other qualified professional authorities.	
gypsy moth	4 - 8 (0.06 - 0.12 lb ai/acre)	Apply to early instars (1st, 2nd, or 3rd) at first signs of infestation.	

Ground Application: Apply a minimum of 30 gallons per acre by conventional ground equipment. Equipment and spray volume should be calibrated to assure uniform coverage of infested parts of the crop.

Aerial Application: Apply a minimum of 10 gallons per acre. Use of higher carrier volume is recommended for heavy infestations and in situations where thorough coverage is difficult to achieve.

Rotational Crop Restrictions: Crops with registered uses may be replanted at any time. All other crops grown for food or feed may be replanted after 7 days.

Note: Drift and runoff from applications of this product may be hazardous to sensitive aquatic invertebrates in water bodies adjacent to the treatment area. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Intrepid® 2F EPA Reg. No. 62719-442

Control of Lepidoptera Larvae Infesting Bushberries (Subgroup 13-07B¹), Aronia Berry, Buffalo Currant, Chilean Guava, European Barberry, Highbush Cranberry, Honeysuckle, Jostaberry, Juneberry, Lingonberry, Native Currant, Salal, Sea Buckthorn, and Cultivars and/or Hybrids of Each

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Intrepid® 2F insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Intrepid 2F according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Intrepid 2F.

Directions for Use

¹Bushberries (subgroup 13-07B) including black currant, elderberry, gooseberry, highbush blueberry, huckleberry, lowbush blueberry, red currant

Refer to product label for General Use Precautions, Mixing and Application instructions.

Pests, Application Rates, Application Timing and Restrictions:

Pests	Application Rate (fl oz/acre)	Application Timing	Restrictions
cherry fruitworm cranberry fruitworm	16 (0.25 lb ai/acre)	Apply at initiation of egg laying [approximately 400 Day Degrees (DD) base 50°F] following biofix. Make a second application at 100% petal fall (usually 7 to 14 days following the first application). An additional application (third) no sooner than 7 days following the second application may be required under high pressure or sustained moth flight.	<ul style="list-style-type: none"> • Do not apply more than 64 fl oz of Intrepid 2F (1 lb ai) per acre per calendar year. • Do not make more than 3 applications per acre per calendar year. • Re-treatment Interval: 7 days • Pre-harvest Interval: Do not harvest within 7 days of application. • See Rotational Crop Restrictions below.
armyworm cutworm	8 – 16 (0.12 – 0.25 lb ai/acre)		
gypsy moth	4 – 8 (0.06 – 0.12 lb ai/acre)		

Page 1 of 3

obliquebanded leafroller		Spring (overwintering) generation: Make one or two applications at bloom to petal fall to small larvae when threshold levels occur. Summer generation: Begin applications at peak moth flight (200-300 DD base 43°F) following biofix. An additional application (third) no sooner than 7 days following the second application may be required under high pressure or sustained moth flight.
redbanded leafroller variegated leafroller		For control of other leafrollers, apply at early egg hatch for each generation. Make the first application before webbing and sheltering begins. Make a second application to ensure complete coverage of rapidly expanding fruits or foliage.
spanworm		Apply when first signs of feeding damage appear or when infestations reach threshold levels as defined by cooperative extension service or other qualified professional authorities.
green fruitworm		Apply when larvae are first detected in the clusters or when infestations reach threshold levels as defined by cooperative extension service or other qualified professional authorities.
armyworm cutworm	8 – 16 (0.12 – 0.25 lb ai/acre)	Apply when first signs of feeding damage appear or when infestations reach threshold levels as defined by cooperative extension service or other qualified professional authorities.
gypsy moth	4 – 8 (0.06 – 0.12 lb ai/acre)	Apply to early instars (1st, 2nd, or 3rd) at first signs of infestation.

¹Biofix is defined as first sustained adult catch in pheromone traps, typically five moths in three traps within a 7-day period. Consult state extension specialists or other qualified authorities for specific information regarding number, placement and management of pheromone traps.

Ground Application: Apply a minimum of 30 gallons per acre by conventional ground equipment. Equipment and spray volume should be calibrated to assure uniform coverage of infested parts of the crop.

Aerial Application: Apply a minimum of 10 gallons per acre. Use of higher carrier volume is recommended for heavy infestations and in situations where thorough coverage is difficult to achieve.

Page 2 of 3

Rotational Crop Restrictions: Crops with registered uses may be replanted at any time. All other crops grown for food or feed may be replanted after 7 days.

Note: Drift and runoff from applications of this product may be hazardous to sensitive aquatic invertebrates in water bodies adjacent to the treatment area. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

Product Bulletin



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Intrepid® 2F EPA Reg. No. 62719-442

Special 2(ee) Recommendation¹
(Not for Use in the State of Texas)

Control of Lepidoptera Larvae Infesting Bushberries (Subgroup 13-07B¹), Aronia Berry, Buffalo Currant, Chilean Guava, European Barberry, Highbush Cranberry, Honeysuckle, Jostaberry, Juneberry, Lingonberry, Native Currant, Salal, Sea Buckthorn, and Cultivars and/or Hybrids of Each

ATTENTION

¹This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to the container for Intrepid® 2F insecticide before applying. Carefully follow all precautionary statements and applicable use directions except as specified below.
- Use of Intrepid 2F according to this Section 2(ee) recommendation is subject to all use precautions and limitations imposed by the label affixed to the container for Intrepid 2F.

Directions for Use

¹Bushberries (subgroup 13-07B) including black currant, elderberry, gooseberry, highbush blueberry, huckleberry, lowbush blueberry, red currant

Refer to product label for General Use Precautions, Mixing and Application instructions.

For control of cherry fruitworm, cranberry fruitworm, obliquebanded leafroller, redbanded leafroller, variegated leafroller, spanworm, and green fruitworm, apply 10 to 16 fl oz (0.16 to 0.25 lb ai) per acre.